

## 2.3 Compute Integration

► I think you have enough skill to solve basic integration of higher secondary level. If you don't have prerequisite knowledge then just go through those chapters.

### 2.3.1 Line Integral

[Do It Yourself] 2.60. Let  $P = \int_0^1 \frac{dx}{\sqrt{8-x^2-x^3}}$  which of the following statements is TRUE

(A)  $\sin^{-1}(\frac{1}{2\sqrt{2}}) \leq P \leq \frac{1}{\sqrt{2}} \sin^{-1}(\frac{1}{2})$  (B)  $\frac{1}{\sqrt{2}} \sin^{-1}(\frac{1}{2}) \leq P \leq \sin^{-1}(\frac{1}{2})$

(C)  $\frac{1}{\sqrt{2}} \sin^{-1}(\frac{1}{2\sqrt{2}}) \leq P \leq \sin^{-1}(\frac{1}{2\sqrt{2}})$  (D)  $\sin^{-1}(\frac{1}{2}) \leq P \leq \frac{\sqrt{3}}{2} \sin^{-1}(\frac{1}{2})$ .

[Hint:  $\int_0^1 \frac{dx}{\sqrt{8-x^2}} \leq \int_0^1 \frac{dx}{\sqrt{8-x^2-x^3}} \leq \int_0^1 \frac{dx}{\sqrt{8-2x^2}}$ ]

[Do It Yourself] 2.61. Solve:  $\int \sqrt{x^2 - a^2} dx$ ,  $\int \sqrt{x^2 + a^2} dx$ ,  $\int \frac{1}{\sqrt{x^2 - a^2}} dx$ ,  $\int \frac{1}{\sqrt{x^2 + a^2}} dx$ ,  $\int \frac{1}{x^2 - a^2} dx$ ,  $\int \frac{1}{x^2 + a^2} dx$ ,  $\int \ln x dx$ ,  $\int \ln(x+1) dx$ ,  $\int \log x dx$ .

[Do It Yourself] 2.62. Solve:  $\int \frac{1}{(x-1)(x-2)(x-3)} dx$ ,  $\int \frac{2x+3}{x^2+1} dx$ ,  $\int \frac{1}{\sin^2 x} dx$ ,  $\int \sin^3 x dx$ ,  $\int \cos^3 x dx$ ,  $\int \sin^4 x dx$ ,  $\int \cos^4 x dx$ .

[Do It Yourself] 2.63. Solve:  $\int_{-1}^2 [x] dx$ ,  $\int_{-1}^2 [x^2] dx$ ,  $\int_{-1}^2 [x]^2 dx$ ,  $\int_{-1}^2 [x+1] dx$ ,  $\int_{-1}^2 |x| dx$ .

Example 2.10. Evaluate:  $\lim_{n \rightarrow \infty} \left[ \frac{1}{n^2 + 1^2} + \frac{2}{n^2 + 2^2} + \frac{3}{n^2 + 3^2} + \cdots + \frac{1}{2n} \right]$ .

$$\begin{aligned} &\Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n} \left[ \frac{n}{n^2 + 1^2} + \frac{2n}{n^2 + 2^2} + \frac{3n}{n^2 + 3^2} + \cdots + \frac{n \cdot n}{n^2 + n^2} \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \left[ \frac{\frac{1}{n}}{1 + (\frac{1}{n})^2} + \frac{\frac{2}{n}}{1 + (\frac{2}{n})^2} + \frac{\frac{3}{n}}{1 + (\frac{3}{n})^2} + \cdots + \frac{\frac{n}{n}}{1 + (\frac{n}{n})^2} \right] = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=1}^n \frac{\frac{r}{n}}{1 + (\frac{r}{n})^2} \\ &= \int_0^1 \frac{x}{1+x^2} dx = \left[ \frac{1}{2} \ln(1+x^2) \right]_0^1 = \frac{1}{2} \ln 2. \end{aligned}$$

[Do It Yourself] 2.64. Find the following limits:

1.  $\lim_{n \rightarrow \infty} \left[ \frac{1^2}{n^3 + 1^3} + \frac{2^2}{n^3 + 2^3} + \cdots + \frac{1}{2n} \right]$ .
2.  $\lim_{n \rightarrow \infty} \left[ \frac{1}{\sqrt{2n-1^2}} + \frac{1}{\sqrt{4n-2^2}} + \cdots + \frac{1}{n} \right]$ .

## 2.3.2 Double Integral

► Repeated Integrals:  $\int_a^b \left[ \int_{\phi_1(x)}^{\phi_2(x)} f(x, y) dy \right] dx$  or,  $\int_c^d \left[ \int_{\psi_1(y)}^{\psi_2(y)} g(x, y) dx \right] dy$ .

► If the inner functions are easily integrable then evaluate using repeated integral else change the order and integrate.

► If the above two methods does not work then we will use the method of transformation.

**Example 2.11.** Find the double integral  $\int_0^9 \int_{\sqrt{x}}^3 \frac{1}{1+y^3} dy dx$ .

$$\Rightarrow \text{Here } I = \int_{x=0}^9 \int_{y=\sqrt{x}}^3 \frac{1}{1+y^3} dy dx = \int_{y=0}^3 \int_{x=0}^{y^2} \frac{1}{1+y^3} dx dy = \int_{y=0}^3 \frac{y^2}{1+y^3} dy = \frac{\ln 28}{3}.$$

**Example 2.12.** What is the value of  $\int_0^{\frac{\pi}{2}} \int_0^x e^{\sin y} \sin x dy dx$ ?

$$\Rightarrow \text{Here } I = \int_{x=0}^{\frac{\pi}{2}} \int_{y=0}^x e^{\sin y} \sin x dy dx = \int_{y=0}^{\frac{\pi}{2}} \int_{x=y}^{\pi/2} e^{\sin y} \sin x dx dy = \int_{y=0}^{\pi/2} e^{\sin y} \left[ -\cos \frac{\pi}{2} + \cos y \right] dy = \int_{y=0}^{\pi/2} e^{\sin y} \cos y dy = e - 1.$$


**Example 2.13.** If  $\int_0^1 \int_y^{2-\sqrt{1-(y-1)^2}} f(x, y) dx dy = \int_0^1 \int_0^{\alpha(x)} f(x, y) dy dx + \int_1^2 \int_0^{\beta(x)} f(x, y) dy dx$

then  $\alpha(x)$  and  $\beta(x)$  are

(A)  $\alpha(x) = x, \beta(x) = 1 + \sqrt{1 - (x-2)^2}$  (B)  $\alpha(x) = x, \beta(x) = 1 - \sqrt{1 - (x-2)^2}$   
 (C)  $\alpha(x) = 1 + \sqrt{1 - (x-2)^2}, \beta(x) = x$  (D)  $\alpha(x) = 1 - \sqrt{1 - (x-2)^2}, \beta(x) = x$ .

$$\Rightarrow \text{Here } I = \int_{y=0}^1 \int_{x=y}^{2-\sqrt{1-(y-1)^2}} f(x, y) dx dy.$$

Now we will draw the area enclosed by the given limits.



Therefore, the option B is correct.

Now we draw  $y = 0, y = 1$ .

$x = y, x = 2 - \sqrt{1 - (y-1)^2}$ .

These four curve generates OABO.

Now we will change the order.

Given  $y$  is constant from 0 to 1.

Changing order we get  $x$  is constant from  $x = 0$  to  $x = 2$ .

First Part:  $x = 0, x = 1$  and  $y = 0, y = x$

Second Part:  $x = 1, x = 2$  and  $y = 0, y = 1 - \sqrt{1 - (x-2)^2}$ .

[Do It Yourself] 2.65. The integral  $\int_0^1 \int_{x^2}^{2x} f(x, y) dy dx$  is equal to

(A)  $\int_0^1 \int_{y/2}^{\sqrt{y}} f(x, y) dx dy + \int_1^2 \int_{y/2}^1 f(x, y) dx dy$  (B)  $\int_0^2 \int_y^{y/2} f(x, y) dx dy$   
(C)  $\int_0^1 \int_{y/2}^{\sqrt{y}} f(x, y) dx dy + \int_1^2 \int_y^{2y} f(x, y) dx dy$  (D)  $\int_0^2 \int_y^{2\sqrt{y}} f(x, y) dx dy$ .

[Do It Yourself] 2.66. Let  $I = \int_0^1 \int_{x^2}^{2-x} xy dy dx$ . The change of order of integration in the integral gives  $I$  as

(A)  $I = \int_0^1 \int_0^{\sqrt{y}} xy dx dy + \int_1^2 \int_0^{2-y} xy dx dy$  (B)  $I = \int_0^1 \int_0^{2-y} xy dx dy + \int_1^2 \int_0^{2-y} xy dx dy$   
(C)  $I = \int_0^1 \int_0^{\sqrt{y}} xy dx dy + \int_0^1 \int_0^{2-y} xy dx dy$  (D)  $I = \int_0^1 \int_0^{2-y} xy dx dy + \int_1^2 \int_0^{\sqrt{y}} xy dx dy$ .

[Do It Yourself] 2.69. Evaluate  $\int_0^1 \int_y^1 x^2 e^{xy} dx dy$ .

[Do It Yourself] 2.70. Evaluate the integral  $\int \int_R \sin(x+y) dx dy$  Where  $R : \{0 \leq x \leq \frac{\pi}{2}; 0 \leq y \leq \frac{\pi}{2}\}$ .

[Do It Yourself] 2.71. Evaluate the integral  $\int \int_R (4 - x^2 - y^2) dx dy$  Where  $R$  is the region bounded by  $x = 0$ ,  $x = 1$ ,  $y = 0$ ,  $y = 3/2$ .

[Do It Yourself] 2.72. Evaluate the integral  $\int \int_R \frac{dx dy}{\sqrt{x^2 + y^2}}$  Where  $R$  is the region bounded by  $|x| \leq 1$ ,  $|y| \leq 1$ .

[Do It Yourself] 2.73. Change the order of integration  $\int_0^1 dx \int_{x^2}^{\sqrt{x}} f(x, y) dy$ .

[Do It Yourself] 2.74. Evaluate  $\int_0^1 \int_y^1 e^{x^2} dx dy$ .

[Do It Yourself] 2.75. Show that  $\int \int_R e^{y/x} dx dy$  Where  $R$  is the triangle bounded by  $y = x$ ,  $y = 0$ ,  $x = 1$  is  $\frac{e-1}{2}$ .

## 2.4 Transformation of variables

► Let  $f$  be a bounded function of  $x$  and  $y$  over a closed region  $S$ . If the transformation  $x = \phi(u, v)$ ,  $y = \psi(u, v)$  represents a continuous bijection between the closed region  $S$  of the  $xy$ -plane onto a region  $S_1$  on  $uv$ -plane and if the functions  $\phi, \psi$  have continuous first order partial derivatives then

$$\int \int_S f(x, y) dx dy = \int \int_{S_1} f[\phi(u, v), \psi(u, v)] \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv.$$

► Here  $J = \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}$  is the Jacobian and is non-zero for transformation.

**Example 2.14.** The value of  $\frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^y e^{-\frac{1}{2}(x^2+y^2)} dx dy$  is

(A)  $\frac{\pi}{4}$  (B)  $\frac{1}{2\pi}$  (C)  $\frac{1}{4}$  (D)  $\frac{1}{2}$ .

⇒ Let  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

Therefore,  $J = \frac{\partial(x, y)}{\partial(r, \theta)} = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = r$ .

$$\begin{aligned} \text{So, } \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^y e^{-\frac{1}{2}(x^2+y^2)} dx dy &= \frac{1}{2\pi} \int_{r=0}^{\infty} \int_{\theta=\pi/4}^{5\pi/4} e^{-\frac{r^2}{2}} r dr d\theta \\ &= \frac{1}{2\pi} \left[ \int_{r=0}^{\infty} e^{-\frac{r^2}{2}} r dr \right] \left[ \int_{\theta=\pi/4}^{5\pi/4} d\theta \right] = \frac{1}{2\pi} \times 1 \times \pi = \frac{1}{2}. \end{aligned}$$

### 2.4.1 Surface Area using Integral

**Example 2.15.** Let  $S = \{(x, y) \in \mathbb{R}^2 : x, y \geq 0, \sqrt{4 - (x - 2)^2} \leq y \leq \sqrt{9 - (x - 3)^2}\}$  then find the area of  $S$ .

⇒ Draw the area carefully.

Now  $\sqrt{4 - (x - 2)^2} = \sqrt{x(4 - x)} \Rightarrow 0 < x < 4$  and this is an upper half of the circle  $(x - 2)^2 + y^2 = 2^2$ .

Also  $\sqrt{9 - (x - 3)^2} = \sqrt{x(6 - x)} \Rightarrow 0 < x < 6$  and this is an upper half of the circle  $(x - 3)^2 + y^2 = 3^2$ .

Now draw the region and you can find the area is  $5\pi/2$ .

[Do It Yourself] 2.82. Let  $S = \{(x, y) \in \mathbb{R}^2 : |x| + |y| \leq 1\}$  then what is the area of  $S$ ?  
[Hint : Easy]

[Do It Yourself] 2.83. Find the value of the real number  $m$  for which

$$\int_0^1 \int_x^{\sqrt{2-x^2}} (x^2 + y^2) dy dx = \int_{m\pi}^{\frac{\pi}{2}} \int_0^{\sqrt{2}} r^3 dr d\theta.$$

[Ans :  $m = 1/4$ , Transform and find the limits of integral]

[Do It Yourself] 2.84. Let  $D$  be the triangle bounded by the  $y$  axis, the line  $2y = \pi$  and the line  $y = x$ . Then the value of the integral  $\int \int_D \frac{\cos y}{y} dx dy$  is

(A) 1/2 (B) 1 (C) 3/2 (D) 2.

[Hint :  $\int \int_D \frac{\cos y}{y} dx dy = \int_{y=0}^{\pi/2} \int_{x=0}^y \frac{\cos y}{y} dx dy$ ]

[Do It Yourself] 2.88. Using the transformation  $x + y = u$ ,  $y = uv$ , show that

$$\int_0^1 dx \int_0^{1-x} \frac{y}{e^{x+y}} dy = \frac{1}{2}(e - 1).$$

[Hint : Draw the region :  $x = 0, y = 0, x + y = 1$ . Now,  $y = 0 \Rightarrow uv = 0 \Rightarrow u = 0, v = 0$ ;  $x = 0 \Rightarrow uv = u \Rightarrow v = 1$ ;  $x + y = 1 \Rightarrow u = 1$ ;  $0 \leq u \leq 1, 0 \leq v \leq 1$ ]

[Do It Yourself] 2.89. Show that  $\int_0^1 dx \int_0^x \sqrt{x^2 + y^2} dy = \frac{1}{6}(\sqrt{2} + \ln(1 + \sqrt{2}))$ .

[Do It Yourself] 2.90. Show that  $\int \int_R (x^2 + y^2) dx dy$ , where  $E$  is the region bounded by  $xy = 1$ ,  $y = 0$ ,  $y = x$ ,  $x = 2$  is  $47/24$ .

[Do It Yourself] 2.91. Show that  $\int \int_E (x^2 + y^2) dx dy = 6/35$ , where  $E$  is the region bounded by  $y = x^2$ ,  $y^2 = x$ .

[Do It Yourself] 2.94. Find range  $W$  of the transformation variable  $(u, v)$  for the transformation  $x = f_1(u, v)$ ,  $y = f_2(u, v)$  and the range  $R$  of  $(x, y)$  are given below

1.  $R$  is bounded by a triangle with vertices  $(0, 0)$ ,  $(1, 0)$ ,  $(0, 1)$ . Transformation  $x = \frac{v(1-u)}{2}$ ,  $y = \frac{v(1+u)}{2}$ . Draw  $W$  and find the integration limits.
2.  $R$  is bounded by a triangle with vertices  $(0, 0)$ ,  $(1, 0)$ ,  $(0, 1)$ . Transformation  $x = v - u$ ,  $y = v + u$ . Draw  $W$  and find the integration limits.
3.  $R$  is bounded by a triangle with lines  $x = 0$ ,  $y = 0$ ,  $x + y = 1$ . Transformation  $x = uv$ ,  $y = u(1 - v)$ . Draw  $W$  and find the integration limits.

[Do It Yourself] 2.95. Show that  $\int \int_R e^{\frac{y-x}{y+x}} dx dy$  Where  $R$  is the triangle bounded by  $y + x = 1$ ,  $y = 0$ ,  $x = 0$  is  $\frac{1}{4}(e - \frac{1}{e})$ .

[Hint : Transformation  $y - x = 2u$ ,  $y + x = 2v$ ]

[Do It Yourself] 2.96. Show that  $\int_0^1 \int_0^{1-x} e^{\frac{y}{x+y}} dx dy = \frac{1}{2}(e - 1)$ .

[Hint : Transformation  $x + y = u$ ,  $y = uv$ ]

[Do It Yourself] 2.98. Find  $\int_0^1 \int_0^x \sqrt{x^2 + y^2} dx dy$ .