

Ch: Set & Seq  
part 3 | Absolute value

$$|x| = \begin{cases} x & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -x & \text{if } x < 0 \end{cases}$$

properties:

- (1)  $|x| = \max \{x, -x\}$
- (2)  $-|x| = \min \{x, -x\}$
- (3)  $|x|^2 = x^2$
- (4)  $|xy| = |x||y|$
- (5)  $|x+y| \leq |x| + |y|$
- (6)  $||x| - |y|| \leq |x-y|$

Hint for (6)  
 $x = x - y + y$   
 $|x| \leq |x-y| + |y|$   
 $\Rightarrow |x| - |y| \leq |x-y|$   
Similarly, (H.W.)  
 $|y| - |x| \leq |x-y|$

H.W. ★ Imp

(1) Show that  $\max\{x, y\} = \frac{x+y+|x-y|}{2}$

$$\min\{x, y\} = \frac{x+y-|x-y|}{2}$$

# Sets in $\mathbb{R}$ (continued)

(1) Derived set: The set of all limit pt of a set  $S$  is said to be its derived set and denoted by  $S'$ .



eg: If  $S = (1, 2)$ ,  $S' = [1, 2]$

If  $S = [1, 2]$ ,  $S' = [1, 2]$



(2) Closed set If it contains all its limit pt

i.e., if  $S' \subseteq S$  then  $S$  is closed.  $\sqrt{2}, \sqrt{3}$

H.W (1) Show  $\mathbb{N}$  is closed. (2)  $\mathbb{Q}$  is neither open nor closed

HW (1)  $\mathbb{R}$  is both open & closed

(2)  $\emptyset$  " " " " " "

(3) A set is open  $\Rightarrow$  its complement  
(close) is close (open)

Dense set: A set  $S$  is said to be dense  
in  $B$  if  $B \subset S'$ .

A set  $S$  is said to be dense in itself  
if  $S \subset S'$ .

ex:  $S$  is dense in  $\mathbb{R}$  if  $S' = \mathbb{R}$   
Hence  $\mathbb{Q}$  is dense in  $\mathbb{R}$ .

v. gmp

# Nested Interval Theorem

Consider  $[a, b]$ . If  $\{[a_n, b_n] : n \in \mathbb{N}\}$  be a family of **CLOSED & BOUNDED** nested intervals

then  $\exists$  one and only one point  $x$  such that

$$x \in \bigcap_{n=1}^{\infty} [a_n, b_n]$$

ex Consider  $[-\frac{1}{n}, \frac{1}{n}]$  for  $n=1, 2, 3, \dots$   
 $0 \in \bigcap_{n=1}^{\infty} [-\frac{1}{n}, \frac{1}{n}]$  ( $\bigcap_{n=1}^{\infty} (0, \frac{1}{n}) \rightarrow$  no point)

H.W. (1) Show that, finite union/intersection of open sets is open

[Hint:  $S_1, S_2, \dots, S_n \rightarrow$  individually open sets  
 $\bigcap_{i=1}^n (x) \subset \bigcap_{i=1}^n S_i \Rightarrow \exists N_i(x) \subset S_i$ . Take ~~finite~~ intersection of finite neighbourhood

(2) Show that 0 is a limit pt of  $\left\{ \frac{1}{n} : n \in \mathbb{N} \right\}$

[Hint: Let  $\epsilon > 0$ .  $\exists n \in \mathbb{N} \ni -\epsilon < \frac{1}{n} < \epsilon$   
 $\Rightarrow \frac{1}{n} \in N'(0, \epsilon) \Rightarrow 0$  is a limit pt of  $\left\{ \frac{1}{n} \right\}$

(3) Verify Bolzano-Weierstrass theorem for the following sets in  $\mathbb{R}$  Just show set is bdd, infinite

- (a)  $\left\{ \frac{n}{n+1} : n \in \mathbb{N} \right\}$  (b)  $\left\{ 1 + \frac{1}{n} : n \in \mathbb{N} \right\}$   
(c)  $\left\{ 1 + \frac{(-1)^n}{n^2} : n \in \mathbb{N} \right\}$

H.W. (1) Verify whether the following sets are open / close / none

(a)  $\{x \in \mathbb{R} : 2x^2 - 5x + 2 < 0\}$


(b)  $\{x \in \mathbb{R} : \cos(x) \neq 0\}$

(c)  $\{(0, 1) \setminus \frac{1}{2^n}, n \in \mathbb{N}\}$

(d)  $\{x \in \mathbb{R} : \sin x = 0\}$

(e)  $\{x \in \mathbb{R} : \sin \frac{1}{x} = 0\}$

(f)  $\bigcup_{n=1}^{\infty} I_n$  where  $I_n = \{x \in \mathbb{R} : \frac{1}{4^n} \leq x \leq \frac{1}{2}\}$




17. W.

$\mathbb{R}$

(1) Prove that a finite subset of  $\mathbb{R}$  is  
close set

(2) Show that finite union/intersection of  
close set is close.  $A_1, A_2, \dots, A_n \rightarrow$  closed





# Borel Set

A set that can be obtained from the union, intersection and relative complement of enumerable collection of close & open sets in  $\mathbb{R}$  is known as Borel set.

eg; (v. imp Borel set) Borel set generated by  $[a, b], [a, b), (a, b], (a, b)$   
 $a, b \in \mathbb{R}$



# Cover



Let  $\mathcal{C}$  be an arbitrary collection of sets  $\{A_\alpha : \alpha \in \Lambda\}$ .  $\mathcal{C}$  is said to be a cover of  $S (\subset \mathbb{R})$  if

$$S \subset \bigcup_{\alpha \in \Lambda} A_\alpha$$



\* If  $\{A_\alpha : \alpha \in \Lambda\}$  all open  $\Rightarrow$  open cover

\* If  $\mathcal{C}$  is a cover of  $S$  and a subcollection  $\mathcal{C}'$  of  $\mathcal{C}$  is also a cover of  $S$  then  $\mathcal{C}' =$  **Sub cover**

# Heine - Borel Theorem

Let  $S \subset \mathbb{R}$  is closed and bounded.

Then every open cover of  $S$

has a finite subcover.



Compact set

$S \subset \mathbb{R}$  is compact set if every

open cover of  $S$  has a finite subcover

Hence, a closed and bounded set in  $\mathbb{R}$  is compact

# SEQUENCE

A sequence is a mapping  $f: \mathbb{N} \rightarrow \mathbb{R}$  and denoted by  $\{f(n)\}_{n \in \mathbb{N}}$  or  $\{x_n\}_{n \in \mathbb{N}}$

Here  $f(n)$  or  $x_n = n$ th term of seq

ex (1)  $x_n = n^2 \Rightarrow$  infinite seq of squares

(2)  $x_n = \frac{1}{n(n-1)} \Rightarrow \{x_n\}$  is a seq for  $n > 2$

(3)  $x_n = \tan\left(\frac{n\pi}{2}\right)$  is NOT a seq (not defined for infinitely many  $n$ )

# Bounds of sequence $\{x_n\}$

\*  $\{x_n\}$  bdd above if  $\exists b \in \mathbb{R} \ni x_n \leq b \forall n \in \mathbb{N}$

\*  $\{x_n\}$  bdd below if  $\exists a \in \mathbb{R} \ni x_n \geq a \forall n \in \mathbb{N}$

$a =$  lower bound,  $b =$  upper bound

\*  $\{x_n\}$  is bdd if both  $a, b$  exists

eg:  $\{\frac{1}{n}\}$  bdd below with supremum 1, infimum 0

$\{(-1)^n\}$  bdd seq (inf -1, sup 1)

H.W. Find sup, inf of  $\{x_n\} = \{(-1)^n \frac{n^2}{n^2+2}\}$  [ans: -1 & 1]

# Convergence of a sequence

A seq  $\{x_n\}$  is said to converge to a real number  $l$  (or, equivalently, we say  $\{x_n\}$  has limit  $l$ ) iff for any given  $\epsilon > 0$ ,  $\exists N \in \mathbb{N} \ni |x_n - l| < \epsilon \forall n \geq N$

$\exists$   $\equiv$  there exists  
 $\in$   $\equiv$  inclusion  
 $\Rightarrow$   $\equiv$  such that  
 $\forall$   $\equiv$  for all

If  $\{x_n\}$  converges to  $l$   
we write  
 $\lim_{n \rightarrow \infty} x_n = l$  or  $x_n \rightarrow l$

Ex (1) Show that,  $\left\{\frac{1}{n}\right\} \rightarrow 0$

Here  $\left|\frac{1}{n} - 0\right| = \frac{1}{n}$ , for any  $\varepsilon > 0$ ,

$\left|\frac{1}{n} - 0\right| < \varepsilon$  if  $n > \frac{1}{\varepsilon}$ . Take  $N = \left[\frac{1}{\varepsilon}\right] + 1$

$\Rightarrow \exists N \in \mathbb{N} \ni \left|\frac{1}{n} - 0\right| < \varepsilon \forall n \geq N$  (proved)

(2) Show that  $\{x^n\} \rightarrow 0$  if  $|x| < 1$

Here  $|x| < 1 \Rightarrow \frac{1}{|x|} = 1 + h$  for some  $h > 0$

Now  $|x^n - 0| = \frac{1}{(1+h)^n} = \frac{1}{1 + nh + \binom{n}{2}h^2 + \dots + h^n} < \frac{1}{nh}$

So,  $\exists \delta > 0$ ,  $|x^n - 0| < \varepsilon$  if  $n > \frac{1}{\delta \varepsilon}$  [Take  $N = \left[\frac{1}{\delta \varepsilon}\right] + 1$ ]

③ Show that :

(i)  $\left\{ \frac{1}{n^p} \right\}$

converges to 0 if  $p > 0$

$(5n+1)^{\frac{1}{4n}}$   
 $(\star+1)^{\frac{1}{4\star}}$   
 $\frac{5}{4} = 1 \frac{1}{4}$

(ii)  $\left\{ (n+1)^{\frac{1}{n}} \right\}$

converges to 1 as  $\star \rightarrow \infty$

(iii)  $\frac{n}{n+1} \rightarrow 1$

(iv)  $\frac{n^2}{n^2+1} \rightarrow 1$

$\left(1 + \frac{1}{n}\right)^n \rightarrow e$   
 $\left(1 + \frac{1}{5n}\right)^{4n} \rightarrow e^{\frac{4}{5}}$

(v) Show that, a constant seq<sup>n</sup> is conv.



~~Q4~~ Show that,  $\{n^{\frac{1}{n}}\}$  conv. to 1  
 (here  $x_n > 0$  for  $n > 1$ )

Here,  $n^{\frac{1}{n}} > 1$  for  $n > 1$   
 $\Rightarrow n^{\frac{1}{n}} = 1 + x_n$   
 We will show  $x_n \rightarrow 0$

$(1+x_n)^n = n$   
 $\Rightarrow n = 1 + n x_n + \binom{n}{2} x_n^2 + \binom{n}{3} x_n^3 + \dots + x_n^n$   
 $\Rightarrow n > \frac{n(n-1)}{2} x_n^2 \Rightarrow x_n < \sqrt{\frac{2}{n-1}}$   
 $\Rightarrow \forall \epsilon > 0, \exists N > 0$  if  $\sqrt{\frac{2}{n-1}} < \epsilon$  ie  $n > \left[ \frac{2}{\epsilon^2} + 1 \right]$

Take  $N$  accordingly

# Divergent and oscillatory seq


$\{x_n\}$  is said to diverge to  $\infty$  if corresponding to any preassigned positive number  $M$ ,  
 $\exists k \in \mathbb{N} \Rightarrow x_n > M \quad \forall n \geq k$ .

$\{x_n\}$  diverge to  $\infty \Rightarrow \lim_{n \rightarrow \infty} x_n = \infty$

\* Not convergent  $\Rightarrow$  divergent.

This type

of seq called oscillatory



# Oscillate finitely

if  $\exists \alpha \ni |x_n| < \alpha \quad \forall n \in \mathbb{N}$


otherwise, it is said to oscillate infinitely.

H.W. Show that  $[n^2 > M \Rightarrow n > \sqrt{M}]$   
 $\{n^2\} \rightarrow \infty$  [Take  $k = [\sqrt{M}] + 1$ ]

$\{(-1)^n\}$  oscillates finitely  
 $\{(-1)^n n\}$  oscillates infinitely

# Theorem

convergent seq has unique limit.



proof: Let  $\{x_n\}$  has two limits  $l_1, l_2$ .

Consider  $\epsilon = \frac{1}{2} |l_1 - l_2| > 0$  [ie,  $|l_1 - l_2| = 2\epsilon$ ]

$\{x_n\} \rightarrow l_1 \Rightarrow \forall \epsilon > 0, \exists n_1 \in \mathbb{N}, \exists |x_n - l_1| < \epsilon \forall n \geq n_1$

$\{x_n\} \rightarrow l_2 \Rightarrow \forall \epsilon > 0, \exists n_2 \in \mathbb{N}, \exists |x_n - l_2| < \epsilon \forall n \geq n_2$

Now,  $|l_1 - l_2| = |l_1 - x_n + x_n - l_2| \leq |x_n - l_1| + |x_n - l_2|$

Take  $n_3 = \max(n_1, n_2) \Rightarrow |l_1 - l_2| < 2\epsilon \forall n \geq n_3$ , Contradiction!